

ABOUT THE AUTHORS



NETAN CHOUDHRY, MD, FRCSC, DABO: Dr. Netan Choudhry is the co-founder and medical director of the Vitreous Retina Macula Specialists of Toronto. He is an internationally recognized Vitreoretinal Surgeon with affiliations at the University of Toronto, Harvard Medical School and Cleveland Clinic Canada. He is universally published and is recognized as a thought leader in retinal imaging and the diagnosis and treatment of rare disorders of the retina and vitreous.



SAMANTHA ORR, MD, BMSc: Dr. Samantha Orr is a recent medical graduate from the Schulich School of Medicine & Dentistry (Class of 2022), currently completing a retina research fellowship with Dr. Netan Choudhry. She has an interest in the quickly evolving clinical utility of ophthalmic imaging, inspired by her past experience as an ophthalmic technician.



MOHAMMAD ALI KHAN, MSc: Mohammad Ali Khan is a medical student at McMaster University (Class of 2023) with a MSc in Mathematics & Statistics. He has an interest in the applications of artificial intelligence in ophthalmic imaging modalities, disease pathophysiology, and prognosis.

Clinical utility of OCTA

Samantha Orr, MD, BMSc, Mohammad Ali Khan, MSc,
Netan Choudhry, MD, FRCSC, DABO

INTRODUCTION

Optical coherence tomography angiography (OCTA) is a novel technology that can produce images of blood flow with unprecedented resolution of all the vascular layers of the retina and choroid in a rapid, non-invasive fashion. The technology dates back to 2005, when researchers demonstrated that blood flow could be visualized using swept-source OCT (SS-OCT) based on properties of the Doppler signal.¹ Since then, advancements have led to the technology becoming widely available for the clinical community, aiding in patient diagnosis and monitoring. This review article will highlight examples of ophthalmic diseases where OCTA has important clinical utility; specifically, diabetic retinopathy, age-related macular degeneration, retinal vein occlusions, white-dot syndromes, and early research into neurodegenerative diseases.

DIABETIC RETINOPATHY

In diabetic retinopathy (DR), OCTA allows for diagnosis, monitoring, and may even aid in the early detection of retinal changes in patients with diabetes.²⁻⁴ A study from 2019 used OCTA to compare perfused capillary density (PCD) in diabetic patients with healthy controls. In this study, diabetic patients were further sub-categorized as having non-proliferative DR (NPDR), proliferative DR (PDR), or no DR.² Diabetic patients with no clinical signs of DR had a significantly higher PCD compared to the control group, providing objective measurement of preclinical retinal vascular changes. The researchers hypothesized that this increase in PCD may have been related to the increased recruitment of capillaries and capillary dilatation.² The NPDR and PDR groups demonstrated progressively decreasing PCD. OCTA also provides notable advantages over the current standard of fluorescein angiography (FA). PDR is characterized by retinal ischemia and the development of neovascularization (NV) at the vitreoretinal interface.² OCTA can measure these retinal NVs via observation of supra-retinal flow signals above the internal limiting membrane (ILM) or with outpouching of the ILM. A study in 2020 compared widefield OCTA to ultra-wide-field FA (UWF-FA) and ultra-wide-field colour fundus photography (UWF-CF) to detect retinal NV in eyes with PDR.³ The study demonstrated that widefield OCTA can identify NV not yet evident on UWF-CF and represents a faster and safer alternative to UWF-FA for surveillance of PDR with comparable diagnostic accuracy. OCTA can also allow for differentiation of subtle NVs from

microaneurysms, which may appear similar on FA.⁴ Another study demonstrated the utility of widefield OCTA through flow overlay on cross-sectional B scans in the staging and prognostication of DR.⁵ Intraretinal microvascular abnormalities (IRMA) are seen on OCTA as collateral vessels within the retina.⁵ As the presence of IRMAs denotes the transition to severe NPDR, widefield OCTA can aid in identifying high-risk DR eyes (**Figure 1**).




Figure 1: Widefield OCTA imaging of diabetic retinopathy seen on OCTA (Zeiss Plex Elite 9000, Oberkochen, Germany). Widefield OCTA of the superior vascular plexus (A) showing areas of non-perfusion, vessel pruning, pre-retinal hemorrhage, and development of neovascularization. Corresponding flow B scan (B) showing subretinal and intraretinal fluid.

NONEXUDATIVE NEOVASCULAR AGE-RELATED MACULAR DEGENERATION

In recent years, widespread clinical use of OCTA has allowed further exploration into age-related macular degeneration (AMD) presenting with macular neovascularization (MNV) without macular fluid or leakage on FA (**Figure 2**).^{6,7} This condition, originally described in the 1970s, is termed non-exudative neovascular AMD.^{6,7} Using en-face OCTA correlated with B scan flow overlay, MNV can be reliably detected and monitored (**Figure 2**).⁶ These eyes are at an increased risk of exudation compared to eyes with non-neovascular AMD.^{6,8-10} The estimated incidence of new onset exudation where the fellow eye had exudative AMD was approximately 25% over different follow-up periods ranging from 6–20 months.⁷ Growth of the MNV has been proposed as a possible biomarker to predict conversion to exudative AMD.⁷ However, further longitudinal research is required, as this finding has not been consistently elucidated across all studies.⁷ Although no randomized clinical trials have been conducted regarding the management of nonexudative neovascular AMD, the consensus is that it should not be treated, but rather monitored.^{6,7} Some studies have proposed that this

neovascularization may have a protective effect against geographic atrophy (GA), with results from some demonstrating a lower rate of GA lesion growth and surviving retinal pigment epithelium over areas of MNV with adjacent atrophy.^{6,7,9} OCT and OCTA technology allows ophthalmologists to closely monitor these patients for the onset of exudation requiring early initiation of treatment.

RETINAL VEIN OCCLUSIONS

In retinal vein occlusions (RVO), OCTA has become particularly valuable in its ability to delineate specific microvascular details in the superficial and deep retinal plexuses and to provide depth-resolved measurements of the foveal avascular zone (FAZ). The size of FAZ measured separately in the superficial and deep vascular plexuses are significantly larger in eyes with RVO compared to fellow eyes, a measurement not possible with the clinical standard using FA.¹¹ Increased FAZ area in the superficial vascular plexus (SVP) has been shown to correlate with poorer visual acuity in RVO.¹¹ Both OCTA and FA can evaluate areas of central and peripheral non-perfusion. However, using OCTA to improve determination of the extent and severity of non-perfusion has important prognostic




Figure 2: Nonexudative neovascular age-related macular degeneration seen on OCTA (Topcon DRI Triton, Tokyo, Japan). Cross-sectional B scan with flow overlay (A) showing pigment epithelial detachment without subretinal or intraretinal fluid. OCTA of the outer retina (B) reveals choroidal neovascularization. Perfusion density mapping of the outer retina (C) highlights area of neovascularization. Follow up images after development of exudation and treatment, including B scan with flow overlay (D), OCTA of outer retina (E), and perfusion density map (F).

implications. Although macular edema is often responsible for vision loss in RVO, photoreceptor damage from non-perfusion can cause persistently poor vision.¹² Specifically, grade 4 macular ischemia, according to the Bradley classification, is correlated with poor visual outcomes.¹² However, grades 2 and 3 macular ischemia show no such significant correlation.¹² OCTA also allows for visualization of microvascular changes including vascular tortuosity, telangiectasia, and collateral vessel development.¹³

WHITE DOT SYNDROMES

The use of OCTA in white dot syndromes has provided insight into disease pathophysiology, diagnosis, and management.¹⁴⁻¹⁷ Although these syndromes are rare and can be clinically similar, OCTA has allowed further characterization and differentiation of these pathologies. Lesions in multiple evanescent white dot syndrome (MEWDS) and acute posterior multifocal placoid pigment epitheliopathy (APMPPE) have clinically similar features.¹⁴⁻¹⁸ Use of OCT and OCTA technology demonstrates distinct pathophysiologic processes for these conditions. Normal OCTA of the choriocapillaris in MEWDS has shown that this is likely not a choriocapillaris pathology,

but rather a primary photoreceptor inflammatory process.¹⁵ In APMPPE, OCTA reveals patchy flow deficits in the choriocapillaris (**Figure 3**) co-localized with inflammatory lesions, suggesting primary choriocapillaris ischemia with a secondary effect on the outer retinal layers and the retinal pigment epithelium.¹⁶ Although both MEWDS and APMPPE are generally self-limiting, it is important to distinguish between these conditions. MEWDS can be a masquerade for multifocal choroiditis, syphilis, and vitreoretinal lymphoma.¹⁴ Rarely, APMPPE can be complicated by central nervous system vasculitis.¹⁴ The utility of OCTA in detecting choroidal neovascularization is well known.¹⁷ This is useful in white dot syndromes such as birdshot chorioretinopathy and idiopathic multifocal chorioiditis (iMFC) with panuveitis, which can develop neovascularization secondary to ischemia, requiring treatment.^{14,17} Choriocapillaris flow deficits or reduced flow are seen on OCTA in other white dot syndromes, such as acute idiopathic maculopathy (AIM) and serpiginous or serpiginous-like chorioiditis.^{14,17} Conversely, acute macular neuroretinopathy (AMN) is associated with deep capillary plexus ischemia.¹⁴




Figure 3: Acute posterior multifocal placoid pigment epitheliopathy and multiple evanescent white dot syndrome seen on OCTA (Topcon DRI Triton, Tokyo, Japan). APMPPE: Single cross-sectional B scan through the fovea (A) demonstrates increased hyperreflectivity in the outer nuclear layer and RPE mottling. 6x6 OCTA of the choriocapillaris (C) shows patchy areas of non-perfusion. MEWDS: Single cross-sectional B scan through the fovea (B) shows focal areas with disruption of the ellipsoid zone. 6x6 OCTA of the choriocapillaris (D) shows normal vasculature with no patchy areas of non-perfusion.

NEURODEGENERATIVE DISEASES

The use of OCTA in patients with neurodegenerative conditions such as Alzheimer's dementia (AD) is an evolving area of research.¹⁹⁻²³ Given the similarities between the microvasculature in the retina and brain, retinal vascular findings on OCTA may provide diagnostic or predictive value for neurodegenerative conditions.¹⁹ In patients with AD, some studies have reported statistically significant OCTA findings compared with control groups and others have found correlations between OCTA measurements and cognitive scores measured by tests such as the Montreal Cognitive Assessment (MoCA).^{20,22,24,25} In the eyes of these patients, significantly decreased vessel density in the SVP has been reported.²⁰ Some studies have also reported a significant decrease in the deep vascular plexus (DVP) density and radial peripapillary capillary level as well, although this has not been reliably replicated.²⁰⁻²² A machine learning algorithm used multimodal retinal imaging and patient information to detect AD with reasonable success (AUC of 0.841 in the best-performing model).²⁵ The algorithm paid particular attention to FAZ size along with SVP density to help make the differentiation.²⁵ In patients with no diagnosis of cognitive impairment or dementia, density in the SVP was found to correlate with cognitive test scores.¹⁹ These findings could play a role in early detection of cognitive changes.¹⁹ Further research with standardized OCTA acquisition and longitudinal approaches is required to elucidate the full potential of OCTA in the field of cognitive decline.^{20,22}

CONCLUSION

OCTA is establishing itself as a valuable tool which provides clinicians with useful diagnostic information in many retinal diseases alongside information that can aid disease management. Furthermore, its use extends beyond the retina and provides potential insights into other ocular conditions and even systemic disease. The evidence for OCTA in clinical practice is growing, with new studies published each year, revealing important insights into the nature and treatment of disease. OCTA provides advantages over the current standard of ophthalmic vascular imaging, allowing for rapid acquisition of non-invasive and reproducible images. Despite these advantages, OCTA-based clinical evidence is still in the early stages of development and further clinical trials are needed to allow for the implementation of this imaging modality into guidelines for various ophthalmic diseases. The standardization of imaging protocols and artifact management will be central to a more widespread adoption of this emerging technology.

References

1. Spaide RF, Fujimoto JG, Waheed NK, Sadda SR, Staurengli G. Optical coherence tomography angiography. *Prog Retin Eye Res*. 2018;64:1. doi:10.1016/j.preteyeres.2017.11.003
2. Rosen RB, Andrade Romo JS, Krawitz BD, et al. Earliest Evidence of Preclinical Diabetic Retinopathy Revealed Using Optical Coherence Tomography Angiography Perfused Capillary Density. *Am J Ophthalmol*. 2019;203:103-115. doi:10.1016/j.ajo.2019.01.012
3. Pichi F, Smith SD, Abboud EB, et al. Wide-field optical coherence tomography angiography for the detection of proliferative diabetic retinopathy. *Graefes Arch Clin Exp Ophthalmol*. 2020;258:1901-1909. doi:10.1007/s00417-020-04773-x
4. Hwang TS, Jia Y, Gao SS, et al. Optical Coherence Tomography Angiography Features of Diabetic Retinopathy. *Retina*. 2015;35(11):2371-2376. doi:10.1097/IAE.0000000000000716
5. Arya M, Sorour O, Chaudhri J, et al. Distinguishing Intraretinal Microvascular Abnormalities from Retinal Neovascularization using Optical Coherence Tomography Angiography. *Retin J Retin Vitr Dis*. 2019;00(00):1-10. doi:10.1097/IAE.0000000000000267
6. De Oliveira Dias JR, Zhang Q, Garcia JMB, et al. Natural History of Subclinical Neovascularization in Nonexudative Age-Related Macular Degeneration Using Swept-Source OCT Angiography. *Ophthalmology*. 2018;125(2):255-266. doi:10.1016/j.ophtha.2017.08.030
7. Laiginhas R, Yang J, Rosenfeld PJ, Falcão M. Non-Exudative Macular Neovascularization – A systematic review of prevalence, natural history, and recent insights from OCT angiography. *Ophthalmol Retin*. 2020;4(7):651-661. doi:10.1016/j.oret.2020.02.016
8. Yanagi Y, Mohla A, Lee SY, et al. Incidence of Fellow Eye Involvement in Patients With Unilateral Exudative Age-Related Macular Degeneration. *JAMA Ophthalmol*. 2018;136(8):905. doi:10.1001/JAMAOPHTHALMOL.2018.2154
9. Capuano V, Miere A, Querques L, et al. Treatment-Naive Quiescent Choroidal Neovascularization in Geographic Atrophy Secondary to Nonexudative Age-Related Macular Degeneration. *Am J Ophthalmol*. 2017;182:45-55. doi:10.1016/j.ajo.2017.07.009
10. Heiferman MJ, Fawzi AA. Progression of subclinical choroidal neovascularization in age-related macular degeneration. *PLoS One*. 2019;14(6). doi:10.1371/JOURNAL.PONE.0217805
11. Ouederni M, Khalifa MBH, Sassi H, Nefaa F, Ayed O, Cheour M. Quantitative Analysis of Microvascular Network with Optical Coherence Tomography Angiography and its Correlation with Visual Acuity in Retinal Vein Occlusion. *J Curr Ophthalmol*. 2021;33:453-460. doi:10.4103/JOCO.JOCO_163_21
12. Mejia ME, Rios HA, Rosenstiel S, Rodríguez FJ. Optical coherence tomography angiography as predictor of visual outcomes in retinal vein occlusion treated with antiangiogenic therapy. *Eur J Ophthalmol*. 2022;0(0):1-7. doi:10.1177/11206721221099487
13. Tsai G, Banaee T, Conti FF, Singh RP. Optical coherence tomography angiography in eyes with retinal vein occlusion. *J Ophthalmic Vis Res*. 2018;13(3):315-332. doi:10.4103/jovr.jovr_264_17
14. Pradas M, Rodriguez-Merchante MP, Estébanez N, et al. Navigating the White Dot Syndromes with Optical Coherence Tomography (OCT) and OCT Angiography (OCT-A). *Ocul Immunol Inflamm*. 2022;1-11. doi:10.1080/09273948.2022.2046798
15. Agarwal A, Invernizzi A. The Role of Optical Coherence Tomography and Optical Coherence Tomography Angiography in the Differential Diagnosis of Posterior Uveitis. *Ocul Immunol Inflamm*. 2022;1-8. doi:10.1080/09273948.2022.2071743
16. Testi I, Modugno RL, Pavesio C. Multimodal imaging supporting the pathophysiology of white dot syndromes. *J Ophthalmic Inflamm Infect*. 2021;11(32):1-7. doi:10.1186/S12348-021-00261-3
17. Pichi F, Sarraf D, Morara M, Mazumdar S, Neri P, Gupta V. Pearls and pitfalls of optical coherence tomography angiography in the multimodal evaluation of uveitis. *J Ophthalmic Inflamm Infect*. 2017;7(20):1-12. doi:10.1186/S12348-017-0138-Z
18. Pichi F, Srivastava SK, Chexal S, et al. En face optical coherence tomography and optical coherence tomography angiography of multiple evanescent white dot syndrome: New insights into pathogenesis. *Retina*. 2016;36:S178-S188. doi:10.1097/IAE.0000000000001255
19. Fang M, Strand K, Zhang J, et al. Retinal vessel density correlates with cognitive function in older adults. *Exp Gerontol*. 2021;152:1-6. doi:10.1016/j.exger.2021.111433
20. Wang X, Zhao Q, Tao R, et al. Decreased Retinal Vascular Density in Alzheimer's Disease (AD) and Mild Cognitive Impairment (MCI): An Optical Coherence Tomography Angiography (OCTA) Study. *Front Aging Neurosci*. 2021;12:1-10. doi:10.3389/fnagi.2020.572484
21. Abraham AG, Guo X, Arsiwala LT, et al. Cognitive decline in older adults: What can we learn from optical coherence tomography (OCT)-based retinal vascular imaging? *J Am Geriatr Soc*. 2021;69:2524-2535. doi:10.1111/JGS.17272
22. Yan Y, Wu X, Wang X, et al. The Retinal Vessel Density Can Reflect Cognitive Function in Patients with Alzheimer's Disease: Evidence from Optical Coherence Tomography Angiography. *J Alzheimer's Dis*. 2021;79:1307-1316. doi:10.3233/JAD-200971
23. Wang X, Wei Q, Wu X, et al. The vessel density of the superficial retinal capillary plexus as a new biomarker in cerebral small vessel disease: an optical coherence tomography angiography study. *Neurol Sci*. 2021;42:3615-3624. doi:10.1007/S10072-021-05038-Z
24. Lee MJ, Abraham AG, Swenor BK, Sharrett AR, Ramulu PY. Application of optical coherence tomography in the detection and classification of cognitive decline. *J Curr Glaucoma Pract*. 2018;12(1):10-18. doi:10.5005/JP-JOURNALS-10028-1238
25. Wisely CE, Wang D, Henao R, et al. Convolutional neural network to identify symptomatic Alzheimer's disease using multimodal retinal imaging. *Br J Ophthalmol*. 2022;106(3):388-395. doi:10.1136/BJOPHTHALMOL-2020-317659